



Liquid Crystals of Dendron-like Pt Complexes - Phase 2

Eduardo Arias Marin
CIQA

11/26/2014
Final Report

DISTRIBUTION A: Distribution approved for public release.

Air Force Research Laboratory
AF Office Of Scientific Research (AFOSR)/ IOS
Arlington, Virginia 22203
Air Force Materiel Command

REPORT DOCUMENTATION PAGE				<i>Form Approved</i> OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</small>					
1. REPORT DATE (DD-MM-YYYY)		2. REPORT TYPE		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area code)

Final Report

**Liquid Crystals of Dendron-Like Pt Complexes
Processable Into Nanofilms. Dendrimers**

Eduardo Arias, Ivana Moggio, Ronald Ziolo

Centro de Investigación en Química Aplicada
Blvd. Enrique Reyna, No. 140
Saltillo, Coahuila, México 25294
eduardo.arias@ciqa.edu.mx

AFOSR FA9550-12-1-0234
August 2014

Cholesteric liquid crystal glass platinum acetylides

Eduardo Arias, Ivana Moggio and Ronald F. Ziolo

Centro de Investigación en Química Aplicada
Saltillo, Coahuila, Mexico

Grant Objectives: no changes.

Status of Effort: all research objectives have been met.

Summary: To prepare cholesteric liquid crystalline nonlinear optical materials with ability to be vitrified on cooling and form long time stability cholesteric glasses at room temperature, a series of platinum acetylide complexes modified with cholesterol has been synthesized. The materials synthesized have the formula *trans*-Pt(PR₃)(cholesterol (3 or 4)-ethynyl benzoate)(1-ethynyl-4-**X**-benzene), where **R** = **Et**, **Bu** or **Oct** and **X** = **H**, **F**, **OCH₃** and **CN**. A cholesteric liquid crystal phase was observed in the complexes **R** = **Et**, and **X** = **F**, **OCH₃** and **CN** but not in any of the other complexes. When **X** = **CN**, a cholesteric glass was observed at room temperature which remained stable up to 130 °C, then converted to a mixed crystalline/cholesteric phase and completely melted to an isotropic phase at 230 °C. When **X** = **F** or **OCH₃** the complexes were crystalline at room temperature with conversion to the cholesteric phase upon heating to 190 and 230 °C, respectively. In the series **X** = **CN**, **OCH₃** and **F**, the cholesteric pitch was determined to be 1.7, 3.4 and 9.0 μ, respectively.

INTRODUCTION


Platinum acetylides are nonlinear optical materials with high linear transmission, broadband triplet state spectra and efficient conversion to the triplet state[1]. We have been investigating the relation between chemical structure and spectroscopic properties of platinum acetylide complexes having the molecular formula *trans*-Pt(PBu₃)₂L₂ including two photon spectroscopy. We have processed these chromophores into solid state optical elements[2] and glass-forming liquids[3]. To increase our understanding of the excited state behavior in a chiral environment, we synthesized platinum acetylides having a cholesteric liquid crystal phase. Cholesteric glasses are useful as large area non-absorbing polarizers[4], optical notch filters and reflectors[5], optically-switchable notch filters[6] and polarizing fluorescent films[7] and one-dimensional photonic band-gap for circularly polarized lasing[8]. In this proceeding we describe the liquid crystal behavior of a series of cholesterol-containing platinum acetylides having the general formula shown in Table 1. We found a combination of **R**, **A**, **B**, **X** and **Y** which yielded a stable cholesteric glassy liquid crystal platinum acetylide. The

necessary conditions for cholesteric phase are **R** = Et, **A** = COO-Cholesterol, **B** = H, **X** = polar group and **Y** = H.

EXPERIMENT

Table 1 lists the compounds we synthesized. The structure variations explored the effect of the length of the **R** group, *meta*- or *para*- cholesterol ester substitution(**A**,**B** and **Y**) and the effect of a para substituent(**X**). The method of thermal polarizing microscopy was used in order to characterize these materials[9]. ITO-glasses were covered with polyimide by spin-coating (20 s; 2,000 rpm), pre-baked (~80 °C, ~3-5 min) and baked (~180 °C, 1 hr). Polyimide layers were rubbed in anti-parallel directions with velvet cloth, then fiber spacers (3-5 μ) were sputtered over the surface of the one (bottom) of two substrates. Second (top) and the bottom substrates were clamped, and sealed with UV-glue (NOA65). An empty cell with powders of material on one of its edges was placed on the hot stage. The hot stage was heated above the melting temperature of the material under study until powders were melted, and the cell was filled due to capillary forces. The time required to fill empty cell was about 10-30 min. Cholesteric pitch was determined using a Cano-Grandjean wedge. Two plates with planar boundary conditions were assembled to form a wedge with an opening angle β . This angle was measured independently and was about 1 mrad. The cholesteric helix is perpendicular to the wedge substrates due to planar boundary conditions. When the cell gap increases continuously along the wedge, the integer number of half pitches increases in a discontinuous way through disclination lines. These lines can be observed experimentally. If S is a distance between two disclination lines, then the following expressions is used to calculate the pitch P : $\frac{P}{2 \cdot S} = \tan \beta$. The main challenge of the experiments was to create alignment for all three liquid crystals as the process of their orientation requires usage of high temperatures. Common materials used in liquid crystal technologies fail at around 200 °C. To align these three liquid crystals we needed to warm our samples till approximately 250 °C and then cool them down slowly to induce the necessary planar alignment. To prepare high temperature cells, we sputtered inorganic SiO_x on a glass surface at oblique incidence. This inorganic material withstood higher temperature than those normally used in these experiments.

Table 1. List of complexes synthesized

					
Compound	A	B	R	X	Y
I	H	COOChol ¹	Bu	H	COOChol
II	H	COOChol	Bu	H	H
III	H	COOChol	Bu	CN	H
IV	COOChol	H	Et	H	H
V	COOChol	H	Et	F	H
VI	COOChol	H	Et	CN	H
VII	COOChol	H	Et	OCH ₃	H
VIII	H	COOChol	Oct	H	COOChol
IX	COOQ ²	H	Et	COOQ	H
X	H	COOChol	Et	CN	H
XI	COOChol	H	Bu	CN	H
XII	COOQ	H	Et	CN	H

¹Chol = Cholesterol ²Q = C₅H₁₁COOChol

DISCUSSION

Phase behavior. Material **I** is a crystalline solid which melts to an isotropic phase at 231.2 °C. Material **II** has asymmetry with one *meta*-cholesterol ester and is a crystalline solid which melts to an isotropic phase at 121.6 °C. We added a polar **CN** group in material **III**. It exhibits polycrystalline phase at room temperature. The polycrystalline structure melts at ~ 170 °C to isotropic liquid at 180 °C. On cooling to room temperature it shows isotropic texture. Material **IV** has **R** = **Et** and is a crystalline solid which melts to an isotropic phase 120-200 °C. Material **V**, **X** = **F**, exhibits a room temperature polycrystalline phase. The polycrystalline structure melts at ~ 175-220 °C. On cooling, a cholesteric phase appears in the temperature range 188.2 °C-160 °C (Fig 1A). At 160 °C crystallization takes place. Material **VI**, **X** = **CN**, exhibits a cholesteric phase at room temperature. In the range 130-140 °C, polycrystalline structure appears co-existing with cholesteric phase in the temperature range ~130 – 220 °C. At 228-235 °C, is a transition to isotropic liquid occurs. MDSC measurements show crystallization and melting at 162-210 °C and a glass transition at 162-187 °C. A cholesteric phase appears at ~240 °C down to room temperature(Fig. 1B, Fig. 2). MDSC measurements on cooling show no crystallization signal. The polycrystalline Material **VII**, **X** = **OCH₃**, melts at ~235 °C, and the

melting process to isotropic liquid is completed at 240 °C. On cooling, cholesteric phase appears at ~229 °C and the sample exhibits cholesteric phase in the temperature range 229 °C-188.8 °C (Fig 1C). At 188.8 °C crystallization takes place over 20-30 min.




Figure 1. Selected photographs from thermal polarizing microscopy A: Material **V**, 175 °C, cooling, field of view: 500 x 750 μm ; B: Material **VI**, 25 °C, cooling, field of view: 500 x 750 μm ; C: Material **VII**, 200 °C, cooling, field of view 500 x 750 μm ; D: Material **X**, 22 °C, cooling, field of view: 900 x 1200 μm .




Figure 2. Cholesteric fingerprint texture observed in glassy thin film prepared from compound **VI**, field of view 260 x 340 μm .

Materials **VIII** and **IX** are isotropic liquids at room temperature. Material **X** exhibits a polycrystalline phase at room temperature. When temperature is increased, polycrystalline structure melts at ~160 °C. On cooling to room temperature, the sample under study shows isotropic texture (Fig. 1D). Material **XI** exhibits a polycrystalline phase at room temperature. When the temperature is increased, the

polycrystalline structure melts at ~ 175 - 180 $^{\circ}\text{C}$. On cooling to room temperature the sample shows isotropic texture up to 60 $^{\circ}\text{C}$ and crystallization takes place at reaching 50 $^{\circ}\text{C}$. Material **XII** is an isotropic liquid at room temperature.

Cano-Grandjean wedge. Figure 3 shows photograph of samples used for cholesteric pitch determination. The optical microscope was used to measure the width of Cano - Grandjean stripes for material **VI**. In the case of materials **V** and **VII**, the optical microscope was not used in order to determine cholesteric pitch P due to the large values of P . In this instance, the wedge cell was heated on the hot stage, then it was taken out, placed between two crossed polarizers, and snapshots were taken. The cholesteric pitch P follows the substituent trend $\text{X}=\text{F}(9.0\text{ }\mu) > \text{OCH}_3(3.5\text{ }\mu) > \text{CN}(1.7\text{ }\mu)$. Cholesteric phase of material **VI** was characterized at high temperature (~ 150 $^{\circ}\text{C}$) and at room temperature as the cholesteric phase was preserved at low temperatures as an overcooled state. In both cases the pitch was $1.7\text{ }\mu$. We performed Gaussian 09 DFT calculations(B3LYP/LANL2DZ) on methyl ester analogues of these compounds and found a calculated dipole moment trend $\text{X}=\text{CN}(4.6\text{ D}) > \text{OCH}_3(3.2\text{ D}) > \text{F}(1.8\text{ D})$ which is consistent with the observed cholesteric pitch trend.




Figure 3. A: Photograph of wedge cell ($\beta = 1.45$ mrad) filled with material **VI** cooled to ~ 22 $^{\circ}\text{C}$. B: wedge cell ($\beta = 1.1$ mrad) filled with material **VII**; colorful horizontal stripes are seen in the upper part of the picture($T=230$ $^{\circ}\text{C}$ - 190 $^{\circ}\text{C}$). C: Wedge cell ($\beta = 1.0$ mrad) filled with material **V** on cooling($T=190$ $^{\circ}\text{C}$ - 160 $^{\circ}\text{C}$); colorful horizontal stripes are seen in the upper part of the picture.

Figure 4 summarizes the phase behavior of these compounds. When $\text{X} = \text{H}$, no cholesteric phase is observed, but when $\text{X} =$ polar group(**F**, **OCH₃** or **CN**), a cholesteric phase appears. When $\text{R} = \text{Bu}$ or **Oct**, no cholesteric phase appears, but is observed with $\text{R} = \text{Et}$. Even though $\text{X} = \text{CN}$ and $\text{R} = \text{Et}$ in compound **X**, its bent geometry gives no cholesteric phase, while the linear geometry of compound **VI**

gives a cholesteric phase. The substituent effects on phase behavior are consistent with cholesteric phase appearing in compounds that are chiral, have linear shape, good ability to pack and also the presence of a polar group.





Figure 4. Summary of phase behavior.

DENDRIMERS

New Denrimers. The synthesis, purification and some spectral characteristics of the new dendrimers shown in Fig 5 were reported in AFOSR FA9550-11-1-0169, May, 2013. Further characterizations of the structures are in progress.






Figure 5. New dendrimer structure summary.

CONCLUSIONS

In summary, we have synthesized a series of platinum acetylide cholesterol esters which exhibit cholesteric liquid crystal behavior. Materials **V**, **VI** and **VII** are monotropic: they form cholesteric mesophase only under cooling. In particular, compound **VI** forms an overcooled cholesteric glass at room temperature which remains stable up to 130 °C. Materials **V** and **VII** form a polycrystalline phase at room temperature. We are currently investigating chiral platinum acetylides with higher helical twisting power.

New Discoveries: there are no inventions to be patented as a result of this research grant. The results claimed that report new molecules are important for basic research. However the following full scientific paper is going to be submitted for its publication to *EMRS proceedings, 2014* “Cholesteric liquid crystal glass platinum acetylides,”

REFERENCES

- [1]. (a) J.E. Rogers, J.E. Slagle, D.M. Krein, A.R. Burke, B.C. Hall, A. Fratini, P.A. Fleitz, T.M. Cooper, M. Drobizhev, N.S. Makarov, A. Rebane, K.-Y. Kim, R. Farley and K.S. Schanze, *Inorg. Chem.*, 46, 6483 (2007). (b) J.E. Haley, D.M. Krein, J.L. Monahan, A.R. Burke, D.G.

- McLean, J.E. Slagle, A. Fratini and T.M. Cooper, *J. Phys. Chem. A*, 115, 265 (2011); (c) A. Rebane, M. Drobizhev, N.S. Makarov, E. Beuerman, J.E. Haley, D.M. Krein, A.R. Burke, J.L. Flikkema and T.M. Cooper, *J. Phys. Chem. A*, 115, 4255 (2011). (d) T.M. Cooper, D.M. Krein, A.R. Burke, D.G. McLean, J.E. Haley, J. Slagle, J. Monahan and A. Fratini, *J. Phys. Chem. A*, 116, 139 (2012).
- [2]. C. Liao, A.H. Shelton, K.-Y. Kim and K.S. Schanze *ACS Appl. Mater. Interfaces*, 3, 3225 (2011).
- [3]. J.E. Slagle, T.M. Cooper, D.M. Krein, J.E. Rogers, D.G. McLean and A.M. Urbas, *Chem. Phys. Lett.*, 447, 65 (2007).
- [4]. (a) T.J. Bunning, P.T. Mather, W. Barnes and P.J. Hood, *Liq. Cryst.*, 26, 557 (1999); (b) S.H. Chen, J.C. Mastrangelo and R. J. Jin, *Adv. Matt.*, 11, 1135 (1999).
- [5]. H. Chen, D. Katsis, J. Mastrangelo, S. Chen, S. Jacob and P.J. Hood, *Adv. Matt.* 12, 1283 (2000).
- [6]. N.S. Kumar, S. Abraham, K. Ratheesh, N. Tamaoki, S. Furumi and S. Das, *J. Photochem. Photobiol. A: Chemistry*, 73, 207 (2009).
- [7]. N. Tamaoki N., *Adv. Matt.*, 13, 1135 (2001).
- [8]. S. Furumi, *Chem. Rec.*, 10, 394 (2010).
- [9]. I. Dierking, I., *Textures of Liquid Crystals*, (Wiley-VCH Verlag GmbH & Co., Weinheim, 2003)